



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

KPC-50 confers resistance to ceftazidime-avibactam associated with reduced carbapenemase activity

Poirel, Laurent ; Vuillemin, Xavier ; Juhas, Mario ; Masseron, Amandine ; Bechtel-Grosch, Ursina ; Tiziani, Simon ; Mancini, Stefano ; Nordmann, Patrice

Abstract: KPC-50 is a KPC-3 variant identified from a *Klebsiella pneumoniae* clinical isolate recovered in Switzerland in 2019. As compared to KPC-3, KPC-50 shows i) a three amino-acid insertion (Glu-Ala-Val) between amino acids 276 and 277 amino acid sequence, (ii) an increased affinity to ceftazidime, (iii) a decreased sensitivity to avibactam, explaining the ceftazidime-avibactam resistance, and (iv) associated to a sharp reduction of its carbapenemase activity.

DOI: <https://doi.org/10.1128/AAC.00321-20>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-187914>

Journal Article

Accepted Version

Originally published at:

Poirel, Laurent; Vuillemin, Xavier; Juhas, Mario; Masseron, Amandine; Bechtel-Grosch, Ursina; Tiziani, Simon; Mancini, Stefano; Nordmann, Patrice (2020). KPC-50 confers resistance to ceftazidime-avibactam associated with reduced carbapenemase activity. *Antimicrobial Agents and Chemotherapy*, 64(8):e00321-20.

DOI: <https://doi.org/10.1128/AAC.00321-20>

**KPC-50 confers resistance to ceftazidime-avibactam associated with
reduced carbapenemase activity**

**Laurent Poirel^{a,b,c*}, Xavier Vuillemin^a, Mario Juhas^a, Amandine Masseron^a, Ursina
Bechtel-Grosch^d, Simon Tiziani^e, Stefano Mancini^f, and Patrice Nordmann^{a,b,c,g}**

*^aEmerging Antibiotic Resistance Unit and Medical and Molecular Microbiology, Faculty of
Science and Medicine, University of Fribourg, Fribourg, Switzerland, ^bSwiss National
Reference Center for Emerging Antibiotic Resistance (NARA), University of Fribourg,
Fribourg, Switzerland, ^cINSERM European Unit (IAME, France), University of Fribourg,
Fribourg, Switzerland, ^dDepartment of Trauma Surgery, University Hospital of Zürich,
Zürich, Switzerland, ^eInstitute of Intensive Medicine, University Hospital Zürich, University of
Zürich, Zürich, Switzerland, ^fInstitute of Medical Microbiology, University of Zürich, Zürich,
Switzerland, and ^gInstitut for Microbiology, University Hospital Center and University of
Lausanne, Lausanne, Switzerland*

Running title: KPC-50 variant confers ceftazidime-avibactam resistance

Keywords; KPC, ceftazidime-avibactam, *Klebsiella pneumoniae*

***Address correspondence to:**

Laurent Poirel
Medical and Molecular Microbiology
Faculty of Science and Medicine, University of Fribourg
Chemin du Musée 18

27 CH 1700 Fribourg, Switzerland
28 Phone: +41 26 300 9582
29 E-mail: laurent.poirel@unifr.ch
30

31

32

33 **KPC-50 is a KPC-3 variant identified from a *Klebsiella pneumoniae* clinical**
34 **isolate recovered in Switzerland in 2019. As compared to KPC-3, KPC-50 shows i) a**
35 **three amino-acid insertion (Glu-Ala-Val) between amino acids 276 and 277 amino acid**
36 **sequence, (ii) an increased affinity to ceftazidime, (iii) a decreased sensitivity to**
37 **avibactam, explaining the ceftazidime-avibactam resistance, and (iv) associated to a**
38 **sharp reduction of its carbapenemase activity.**

39

40

41 Occurrence of multidrug-resistant Enterobacterales, and especially of carbapenemase-
42 producing isolates, is increasingly reported, thus leaving very few therapeutic options for
43 treating related infections (1). Interestingly, the recently-marketed ceftazidime/avibactam
44 (CZA) drug combination offers novel interesting perspectives (2). This β -lactam/ β -lactamase
45 inhibitor combo provides a therapeutic alternative for treating infections caused by KPC-like
46 and OXA-48-like producers, whereas producers of carbapenemases of the metallo- β -
47 lactamase type remain resistant to that combination (1, 2). Despite CZA is still rarely
48 prescribed worldwide, KPC-like producing isolates being resistant to this drug combo have
49 been already reported (3-7). Several reports identified KPC variants exhibiting single amino
50 acid substitutions in their omega-loop (amino acid positions 164-179), and particularly the
51 Asp179Tyr substitution, leading to enhanced affinity toward ceftazidime, with a concomitant
52 reduced binding to avibactam (8-12). In addition, we recently identified KPC-41, possessing a
53 three amino-acid insertion in the KPC-3 protein sequence being distantly located from the
54 omega loop (namely between positions 269-270), that conferred high level of resistance to
55 CZA in a clinical *K. pneumoniae* isolate recovered in Switzerland (13).

56

57 *K. pneumoniae* isolate N869 was recovered from a patient repatriated after a traffic
58 accident from Greece to Switzerland. In the Greek hospital, the patient developed a ventilator-
59 associated pneumonia for which he received a treatment made of clindamycin, linezolid and
60 meropenem during two days, to which colistin was added on day 5. Few days later, he was
61 transferred to Switzerland where all antibiotics but meropenem (as monotherapy then) were
62 discontinued. Rectal swabs performed at admission grew *K. pneumoniae* isolate N859 using
63 the Chrom ID Carba Smart[®] selective plate (bioMérieux, La Balme-les-Grottes, France).
64 According to the EUCAST 2020 breakpoints (14), *K. pneumoniae* isolate N859 was resistant
65 to all β -lactams including to imipenem and ertapenem, but remained susceptible to
66 meropenem (Table 1). The carbapenemase activity was evaluated by using the Rapid Carba
67 NP test that gave a positive result (15).

68 *K. pneumoniae* N859 was also resistant to aminoglycosides (kanamycin, tobramycin,
69 netilmicin), to fluoroquinolones and to colistin (MIC at 128 μ g/ml). It remained susceptible to
70 tetracycline, tigecycline, chloramphenicol, trimethoprim-sulfamethoxazole and fosfomycin,
71 and was of intermediate susceptibility to amikacin and gentamicin. It also showed resistance
72 to CZA (MIC at >256 μ g/ml) and to ceftolozane-tazobactam combination (> 256 μ g/ml),
73 using inhibitors concentrations at 4 μ g/ml.

74 PCR identified a *bla*_{KPC}-like gene and sequencing of the corresponding amplicon
75 identified a gene encoding a KPC variant possessing a three amino-acid insertion (Glu-Ala-
76 Val) between amino acids 276 and 277 (Ambler numbering), leading to a novel variant named
77 KPC-50 (Figure S1). Search of additional β -lactamase resistance genes as reported (16)
78 identified a *bla*_{SHV}-like gene (intrinsic to *K. pneumoniae*) but no additional extended-spectrum
79 β -lactamase gene. Mating-out assays performed using *K. pneumoniae* N859 as donor and
80 azide-resistant *Escherichia coli* J53 strain as recipient (13) were successful and confirmed the
81 plasmid location of the *bla*_{KPC-50} gene, being ca. 60-kb in-size (data not shown). No other
82 antibiotic marker was co-transferred along with the *bla*_{KPC-50} gene. PCR-based replicon typing
83 showed that this plasmid belonged to the IncFIB incompatibility group (17). Multilocus
84 sequence typing performed as described (18) showed that isolate N859 belonged to sequence
85 type ST258 that corresponds to the worldwide spread KPC-producing *K. pneumoniae*
86 background (19, 20).

87 In order to confirm if the amino acid substitutions identified within the KPC sequence
88 might be responsible for the CZA resistance phenotype observed in *K. pneumoniae* N859, the
89 *bla*_{KPC-50} gene was cloned and expressed in *E. coli* TOP10. MIC values were then compared
90 with the previously obtained KPC-3-producing *E. coli* TOP10 (13). In such *E. coli*

91 background, KPC-3 conferred resistance to all β -lactams including ceftazidime, but remained
92 susceptible to CZA, as previously shown (13). Conversely, although KPC-50 also conferred
93 high-level resistance to ceftazidime, it additionally conferred high-level of resistance to CZA
94 (Table 1). Noteworthy, KPC-50 conferred much lower level of resistance to ceftazidime,
95 cefotaxime and cefepime than KPC-3. One of the most marked features of KPC-50 as
96 compared to KPC-3 was that its production did not lead to resistance to carbapenems (Table
97 1). Indeed, *E. coli* expressing the *bla*_{KPC-50} gene remained susceptible to ertapenem and
98 meropenem (MICs at 0.25 μ g/ml and 0.5 μ g/ml, respectively), while the MIC of imipenem
99 observed for that KPC-50-producing *E. coli* recombinant strain was at 4 μ g/ml (breakpoint
100 value). Low MICs were observed when testing the new carbapenem / β -lactamase inhibitor
101 combinations such as imipenem/relabactam, and more specifically meropenem/vaborbactam
102 that showed an excellent capacity to inhibit the growth of KPC-50 producers (Table 1).

103 The enzymatic properties of KPC-50 were determined using purified extracts, and
104 compared to those of KPC-3 previously obtained under same conditions (13). Kinetic data
105 showed that KPC-50 has a lower hydrolysis activity of cefalotin, cefotaxime, aztreonam and
106 imipenem, as compared to that of KPC-3 (Table 2). Similar decreased hydrolytic properties
107 toward β -lactams have been previously reported for those KPC variants conferring resistance

108 to the CZA combination, such as for KPC-41 (13) or the Asp179Tyr KPC-2 mutants (21-23).
109 Furthermore, the activity of KPC-50 toward aztreonam was not detectable by contrast to
110 KPC-3 (Table 2), and contrasting also with data obtained for KPC-41 (13).

111 Kinetic activities toward ceftazidime were measured and compared for KPC-50 and
112 KPC-3 enzymes. As expected, a significant hydrolysis rate was detected with KPC-3, but no
113 hydrolysis could be detected with KPC-50 in normal conditions (measurement made during 5
114 min). Then, another assay was performed during 1 hour, showing that ceftazidime was indeed
115 hydrolyzed by KPC-50, but the hydrolysis rate was much lower than that of KPC-3 (Figure).
116 Therefore, we observed here a paradoxical situation with KPC-50 conferring high-level
117 resistance to ceftazidime once produced by a recombinant *E. coli* clone, but weak hydrolysis
118 rate measured by UV spectrophotometry. Thus, compared affinities of KPC-50 and KPC-3 to
119 the ceftazidime substrate were respectively measured using various concentrations of
120 ceftazidime, to inhibit the hydrolysis of a reporter substrate (nitrocefin) as published (13, 24).
121 At the same ceftazidime concentrations, a higher inhibition level of nitrocefin was observed
122 with KPC-50 than with KPC-3 (Figure), showing that KPC-50 exhibited a higher affinity
123 toward ceftazidime compared to KPC-3.

124 Comparative inhibitory activity of clavulanic acid, tazobactam and AVI were
125 determined for KPC-50 and KPC-3, showing a 4-fold lower inhibitory activity of AVI toward
126 KPC-50 compared to KPC-3, while conversely that of tazobactam and clavulanic acid were
127 higher toward KPC-50 than KPC-3 (Table 2).

128 Those results overall indicated that the 276-Glu-Ala-Val-277 insertion observed in the
129 KPC-50 sequence was responsible for a reduced hydrolysis of cefalotin, cefotaxime, and
130 carbapenems, associated to a higher affinity toward ceftazidime and a reduced sensitivity to
131 AVI.

132 CONCLUSION

133 A novel KPC-type enzyme conferring resistance to CZA was identified here from a
134 multidrug-resistant *K. pneumoniae* isolate recovered in Switzerland but likely acquired in
135 Greece, with no known history of treatment with CZA for the patient. Of note, and as already
136 highlighted for KPC-41 and other KPC mutants conferring resistance to CZA, the overall
137 decreased carbapenemase activity observed for KPC-50 might be considered as good news.
138 Furthermore, the newly-developed carbapenem/ β -lactamase inhibitor combinations
139 (meropenem/vaborbactam and imipenem/relebactam) also showed an excellent efficacy

140 against the KPC-50 producers (either the *K. pneumoniae* clinical isolate or the *E. coli*
141 recombinant strain).

142

143 The sequence of KPC-50 has been deposited in the NCBI database under the accession
144 number GenBank MN654342.

145 ACKNOWLEDGMENTS

146 This work was financed by the University of Fribourg, Switzerland, the NARA, and by
147 the Swiss National Science Foundation (projects FNS-31003A_163432 and FNS-
148 407240_177381).

149 AUTHORS CONTRIBUTIONS

150 LP and PN designed the study. RZ and SM provided the material. UBG and ST provided the
151 clinical data. XV, MJ and AM performed the experiments. LP and PN wrote the manuscript.

152 FIGURE LEGEND

153

154 Figure. Analysis of ceftazidime hydrolysis. (A) KPC-50 and KPC-3 (1 μ M enzyme)
155 hydrolysis of 25 μ M ceftazidime (CAZ) at room temperature. (B) and (C) Competitive
156 inhibition curves determined with 50 μ M nitrocefin and increasing concentrations of CAZ

157 with 0.1 μ M KPC-50 (B) and 0.1 μ M KPC-3 (C) at room temperature. For (B) and (C)

158 nitrocefin absorbance was measured.

159

160

161

REFERENCES

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

1. Nordmann P, Poirel L. 2019. Epidemiology and diagnostics of carbapenem resistance in Gram-negative bacteria. Clin Infect Dis 69;S521-S528.
2. Van Duin D, Bonomo RA. 2016. Ceftazidime/avibactam and ceftolozane/tazobactam: second-generation β -lactam/ β -lactamase inhibitor combinations. Clin Infect Dis 63:234-241.
3. Humphries RM, Yang S, Hemarajata P, Ward KW, Hindler JA, Miller SA, Gregson A. 2015. First report of ceftazidime-avibactam resistance in a KPC-3-expressing *Klebsiella pneumoniae* isolate. Antimicrob Agents Chemother 59:6605-6607.
4. Shields RK, Chen L, Cheng S, Chavda KD, Press EG. 2017. Emergence of ceftazidime-avibactam mutations during treatment. Antimicrob Agents Chemother 61:1-11.
5. Hemarajata P, Humphries RM. 2019. Ceftazidime/avibactam resistance associated with L169P mutation in the omega loop of KPC-2. J Antimicrob Chemother;74:1241-1243.
6. Räisänen K, Koivula I, Ilmavirta H, Puranen S, Kallonen T, Lyytikäinen O, Jalava J. 2019. Emergence of ceftazidime-avibactam-resistant *Klebsiella pneumoniae* during treatment, Finland, December 2018. Euro Surveill 24(19).
7. Athans V, Neuner EA, Hassouna H, Richter SS, Keller G, Castanheira M, Brizendine KD, Mathers AJ. 2018. Meropenem-vaborbactam as salvage therapy for ceftazidime-avibactam-resistant *Klebsiella pneumoniae* bacteremia and abscess in a liver transplant recipient. Antimicrob Agents Chemother 63;pii: e01551-18.
8. Barnes MD, Winkler ML, Taracila MA, Page MG, Desarbres E, Kreiswirth BN, Shields RK, Nguyen MH, Clancy C, Spellberg B, Papp-Wallace KM, Bonomo RA.

2017. *Klebsiella pneumoniae* Carbapenemase-2 (KPC-2), substitutions at Ambler
position Asp179, and resistance to ceftazidime-avibactam: unique antibiotic-resistant
phenotypes emerge from β -lactamase protein engineering. *mBio* 8:pii:e00528-17.
9. Livermore DM, Warner M, Jamrozy D, Mushtaq S, Nichols WW, Mustafa N,
Woodford N. 2015. In vitro selection of ceftazidime-avibactam resistance in
Enterobacteriaceae with KPC-3 carbapenemase. *Antimicrob Agents Chemother*
59:5324-5330.
10. Winkler ML, Papp-Wallace KM, Bonomo RA. 2015. Activity of
ceftazidime/avibactam against isogenic strains of *Escherichia coli* containing KPC and
SHV β -lactamases with single amino acid substitutions in the Ω -loop. *J Antimicrob
Chemother* 70:2279-2286.
11. Compain F, Arthur M. 2017. Impaired inhibition by avibactam and resistance to the
ceftazidime-avibactam. *Antimicrob Agents Chemother* 61:1-7.
12. Galani I, Antoniadou A, Karaikos I, Kontopoulou K, Giamarellou H, Souli M. 2019.
Genomic characterization of a KPC-23-producing *Klebsiella pneumoniae* ST258
clinical isolate resistant to ceftazidime-avibactam. 2019. *Clin Microbiol Infect* 25:3-6.
13. Mueller L, Masseron A, Prod'Hom G, Galperine T, Greub G, Poirel L, Nordmann P.
2019. Phenotypic, biochemical and genetic analysis of KPC-41, a KPC-3 variant
conferring resistance to ceftazidime-avibactam and exhibiting reduced carbapenemase
activity. *Antimicrob Agents Chemother* pii: AAC.01111-19.
14. EUCAST. 2020. Breakpoints tables for interpretation of MICs and zone diameters.
Version 10.
http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_10_Breakpoint_Tables.pdf (accessed 01 January 2020)
15. Nordmann P, Poirel L, Dortet L. 2012. Rapid detection of carbapenemase-producing
Enterobacteriaceae. *Emerg Infect Dis* 18:1503-1507.

- 211 16. Girlich D, Naas T, Leelaporn A, Poirel L, Fennewald M, Nordmann P. 2002.
212 Nosocomial spread of the integron-located *veb-1*-like cassette encoding an extended-
213 spectrum β -lactamase in *Pseudomonas aeruginosa* in Thailand. Clin Infect Dis
214 **34**:175-182.
- 215 17. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. 2005.
216 Identification of plasmids by PCR-based replicon typing. J Microbiol Methods
217 63:219-228.
- 218 18. Diancourt L, Passet V, Verhoef J, Grimont PA, Brisse S. 2005. Multilocus sequence
219 typing of *Klebsiella pneumoniae* nosocomial isolates. J Clin Microbiol 43:4178-4182.
- 220 19. Pitout JD, Nordmann P, Poirel L. 2015. Carbapenemase-producing *Klebsiella*
221 *pneumoniae*, a key pathogen set for global nosocomial dominance. Antimicrob Agents
222 Chemother 59:5873-5884.
- 223 20. Munoz-Price LS, Poirel L, Bonomo RA, Schwaber MJ, Daikos GL, Cormican M,
224 Cornaglia G, Garau J, Gniadkowski M, Hayden MK, Kumarasamy K, Livermore DM,
225 Maya JJ, Nordmann P, Patel JB, Paterson DL, Pitout J, Villegas MV, Wang H,
226 Woodford N, Quinn JP. 2013. Clinical epidemiology of the global expansion of
227 *Klebsiella pneumoniae* carbapenemases. Lancet Infect Dis 13:785-796.
- 228 21. Haidar G, Clancy CJ, Shields RK, Hao B, Cheng S, Nguyen MH. 2017. [Mutations in](#)
229 [bla_{KPC-3} that confer ceftazidime-avibactam resistance encode novel KPC-3 variants](#)
230 [that function as extended-spectrum \$\beta\$ -lactamases](#). Antimicrob Agents Chemother
231 61(5).
- 232 22. Shields RK, Nguyen MH, Press EG, Chen L, Kreiswirth BN, Clancy CJ. 2017.
233 Emergence of ceftazidime-avibactam resistance and restoration of carbapenem
234 susceptibility in *Klebsiella pneumoniae* carbapenemase-producing *K. pneumoniae*: a
235 case report and review of literature. Open Forum Infect Dis 4:1-4.

- 236 23. Shields RK, Nguyen MH, Press EG, Chen L, Kreiswirth BN, Clancy CJ. 2017. In
237 vitro selection of meropenem resistance among ceftazidime-avibactam-resistant,
238 meropenem-susceptible *Klebsiella pneumoniae* isolates with variant KPC-3
239 carbapenemases. Antimicrob Agents Chemother 61:pii:e00079-17.
- 240 24. Winkler ML, Papp-Wallace KM, Bonomo RA. 2015. Activity of
241 ceftazidime/avibactam against isogenic strains of *Escherichia coli* containing KPC and
242 SHV β -lactamases with single amino acid substitutions in the Ω -loop. J Antimicrob
243 Chemother 70:2279-2286.
- 244

Table 1. MICs of β -lactams for *K. pneumoniae* clinical isolate N859, *E. coli* TOP10 recombinant strains producing KPC-50 and KPC-3, respectively, and *E. coli* TOP10 recipient strain

β -lactams*	MICs (μ g/ml)			
	<i>K. pneumoniae</i> N859	<i>E. coli</i> TOP10 (pTOPO-KPC-50)	<i>E. coli</i> (pTOPO-KPC-3)	<i>E. coli</i> TOP10
Amoxicillin	>1,024	>256	>128	8
Amoxicillin – clavulanic acid	>1,024	64	>128	8
Ticarcillin	>1,024	>256	>128	8
Ticarcillin – clavulanic acid	>1,024	64	>128	8
Piperacillin	>1,024	256	>128	2
Piperacillin – tazobactam	>1,024	128	>128	2
Cefalotin	>1,024	512	>128	16
Cefotaxime	8	8	>128	<0.125
Cefepime	8	8	128	<0.125
Ceftazidime	2,048	1,024	2,048	0.25
Ceftazidime – avibactam	>256	64	2	0.25
Ceftolozane – tazobactam	>256	64	>256	0.06
Cefoxitin	32	8	128	8
Aztreonam	32	16	64	<0.125
Ertapenem	1	0.25	32	<0.125
Imipenem	16	4	16	<0.125
Imipenem - relabactam	2	0.5	0.5	<0.125
Meropenem	2	0.5	16	<0.125
Meropenem - vaborbactam	<0.125	<0.125	0.5	<0.125

*Clavulanic acid was added at a fixed of 2 μ g/ml, tazobactam at 4 μ g/ml, avibactam at 4 μ g/ml, relabactam at 4 μ g/ml, and vaborbactam at 8 μ g/ml.

Table 2. Kinetic parameters of purified β -lactamases KPC-50 and KPC-3. Data for KPC-3 correspond to those previously reported (13). Inhibitory concentration 50% (IC_{50}) and kinetic inhibition parameters of β -lactamase inhibitors against KPC-50 and KPC-3 (bottom).

β -Lactam(s)*	KPC-50			KPC-3		
	k_{cat} (s ⁻¹)	K_m (μ M)	k_{cat}/K_m (mM ⁻¹ .s ⁻¹)	k_{cat} (s ⁻¹)	K_m (μ M)	k_{cat}/K_m (mM ⁻¹ .s ⁻¹)
Benzylpenicillin	<0.01	ND	ND	5.6	33	0.2
Cefalotin	2.2	30	0.07	47	113.5	0.4
Cefotaxime	1.7	55	0.003	34.9	532.8	0.065
Ceftazidime	<0.01	ND	ND	> 3.3	> 700	> 4.7 E-3
Aztreonam	<0.01	ND	ND	5	194.8	0.03
Imipenem	2	85	0.02	4.7	71.5	0.07
Meropenem	<0.01	ND	ND	0.47	18.5	0.03
Ertapenem	<0.01	ND	ND	0.58	37	0.02

Inhibitor	IC_{50} (μ M)		K_i (μ M)	
	KPC-50	KPC-3	KPC-50	KPC-3
Clavulanic acid	10	20	4	20
Tazobactam	10	50	1	10
Avibactam	4	1	2	1

ND : Not determinable due to a low initial rate of hydrolysis

k_{cat} , turnover ; K_m , Michaelis constant (affinity); k_{cat}/K_m = specificity constant (hydrolysis)

IC_{50} represents the concentration of a drug that is required for 50% inhibition of the enzymatic activity

K_i corresponds to a relative k_{off}/k_{on} to the inhibitor for the enzyme

Figure

